MORPHO
摩芙整形

Post-Operative Medications
术后用药

NAUSEA MEDICATIONS
止吐药物

Scopolamine Patch - Anti-Nausea Patch
- Apply to the skin behind your ear the evening before surgery
- Remove 24 hours after surgery has completed
- This patch helps prevent feeling nauseous from anesthetic medications

东莨菪碱贴剂-止吐贴
- 手术前夜贴于耳后皮肤
- 手术完成后 24 小时取下
- 这种贴剂有助于阻止麻醉药物带来的恶心感
-

Ondansetron 8 mg disintegrating sublingual tablets - Anti-Nausea Medication
- Anti-nausea medication
- Take it every 4 hours as needed for post-operative nausea
- These tablets are placed underneath the tongue and will dissolve and be abdorbed. No need to swallow them, especially when you are already nauseous.

昂旦司琼舌下崩解片（8mg）-止吐药
- 止吐药
- 每 4 小时一次，有效缓解术后呕吐
- 将药片置于舌下，它将自动分解和吸收无需吞咽，尤其是但你已经出现恶心感时

ANTIBIOTICS
抗生素

Keflex - Antibiotic
- Antibiotic taken to prevent post-operative infections
- Take it every 6 hours until you finish all the antibiotics

头孢氨苄 - 抗生素
- 服用抗生素以防止术后感染
- 每 6 小时一次，直至吃完所有抗生素

Cleocin / Clindamycin - Antibiotic

- Antibiotic taken to prevent post-operative infections
- Take it every 6 hours until you finish all the antibiotics

克林霉素 - 抗生素

- 服用抗生素以防止术后感染
- 每 6 小时一次，直至吃完所有抗生素

Tobradex Ophthalmic (Tobramycin/dexamethasone ophthalmic) – Topical Antibiotic

- Topical antibiotic ointment for your eyes
- Ointment: Apply 0.5 inch ointment to conjunctival sac 3 to 4 times a day
- Suspension: Apply 1-2 drops every 4 to 6 hours to each eye

Tobradex 眼膏（妥布霉素/地塞米松眼膏） - 外用抗生素

- 外用抗生素眼膏
- 眼膏：取 0.5 英寸眼膏，涂抹于结膜囊，每天 3-4 次
- 眼药水：每次每眼滴 1-2 滴，每隔 4-6 小时一次

PAIN MEDICATIONS

止痛药

Tylenol #3 (Acetaminophen/Codeine) – Narcotic Pain Medication

- Pain medication for mild to moderately severe pain
- You can take 1-2 tabs every 4 hours as needed for pain

泰诺#3（对乙酰氨基酚/可待因）- 麻醉止痛药

- 用于轻-中度疼痛的止痛药
- 你可每次服用 1-2 片，每隔 4 小时一次，用于止痛

Norco (Acetaminophen/Hydrocodone) – Narcotic Pain Medication

- Pain medication for moderately severe pain
- You can take 1-2 tabs every 4 hours as needed for pain

Norco（对乙酰氨基酚/氢可酮） - 麻醉止痛药

- 用于中度疼痛的止痛药
- 你可每次服用 1-2 片，每隔 4 小时一次，用于止痛

Percocet (Aceetaminophen/Oxycodone) – Narcotic Pain Medication

- Pain medication for moderate to severe pain
- You can take 1-2 tabs every 4 hours as needed for pain
- You can also taper down to ½ tab every four hours if 1 tab is too strong for you
- If you have also been prescribed Flexeril, do not take the Flexeril and Percocet at the same time. Space them out about 2 hours. For example, take Flexeril three time a day, and take Percocet 2 hours after you take the Flexeril, or spaced out during the 8-hour interval between Flexeril doses.

扑热息痛（对乙酰氨基酚/氧可酮） - 麻醉止痛药

- 用于中-重度疼痛的止痛药
- 你可每次服用 1-2 片，每隔 4 小时一次，用于止痛
- 如果每次 1 片的药效太强，你可以减量至每隔 4 小时服用半片
- 如果你的处方里还有环苯扎林（Flexeril），不要同时服用环苯扎林和扑热息痛。两者应间隔约 2 小时。例如，环苯扎林应每日服用三次，你每次服用环苯扎林后，应等待 2 小时或在下次服用前的 8 小时内选择合适时间，再服用扑热息痛。

Neurontin (Gabapentin) – Analgesic for Nerve Pain

Neurontin（加巴喷丁） - 神经痛的止痛药

- 有助于缓解刺痛感，或神经再生引起的针刺痛

MUSCLE RELAXANTS

肌肉松弛剂

Flexeril (Cyclobenzaprine) – Muscle Relaxant

- Muscle Relaxant
- Synergistic with Percocet, and makes Percocet more effective in controlling pain
- Helps alleviate muscle cramping and spasm type pain that other pain medication may not address
- Will decrease the amount of Percocet you need to take to control your pain

Flexeril（环苯扎林） - 肌肉松弛剂

- 肌肉松弛剂
- 与扑热息痛联用，会增加扑热息痛的止痛作用
- 有助于缓解其他止痛药无法解决的肌肉痉挛和痉挛型疼痛
- 会减少你用来控制疼痛的扑热息痛的用量